









RESEARCH ARTICLE

Preliminary genetic barcodes for ash (*Fraxinus*) species and generation of new wide hybrids

William J. Plumb^{1,2,3}  | Laura J. Kelly^{1,2}  | Joe Mullender² | Robyn F. Powell² |
Laszlo Csiba²  | Miguel Nemesio-Gorriz³  | David Carey⁴  |
Mary E. Mason⁴  | William Crowther¹ | Jennifer Koch⁴  | Gerry C. Douglas³ |
Richard J. A. Buggs^{1,2} 

¹School of Biological and Behavioural Sciences, Queen Mary University of London, London, UK

²Science Directorate, Royal Botanic Gardens, Kew, Richmond upon Thames, UK

³Forestry Development Department, Agriculture and Food Development Authority, Teagasc, Dublin, Republic of Ireland

⁴United States Department of Agriculture, Forest Service, Northern Research Station, Delaware, Ohio, USA

Correspondence

Richard J. A. Buggs, School of Biological and Behavioural Sciences, Queen Mary University of London, London, UK.

Email: r.buggs@qmul.ac.uk

Funding information

Erica Waltraud Albrecht Endowment Fund; Department for Environment, Food and Rural Affairs (Defra); Living with Environmental Change Tree Health and Plant Biosecurity Initiative – Phase 2, Grant/Award Number: BB/L012162/1; BBSRC; ESRC; Forestry Commission; NERC; Scottish Government; Teagasc, Grant/Award Number: 2014001

Societal Impact Statement

The world-wide diversity of ash trees includes genetic information encoding resistance to the ash dieback fungus and the emerald ash borer beetle, which are currently devastating ash populations in Europe and North America. In order to mobilise this genetic diversity in conventional breeding programmes, we need to be able to accurately identify ash species from around the world and cross them with one another. Here, we present a preliminary genetic barcoding system for ash, and a series of hybridisation experiments between European ash and other species. Two of the hybrids show early promise against ash dieback.

Summary

- Native ash tree species in Europe and North America are being devastated by ash dieback and the emerald ash borer, respectively. As worldwide ash species differ in their level of susceptibility to these threats, hybrid breeding may allow resistance to be transferred among species. However, we do not know the extent to which distantly related ash species can be crossed, and many ash species are difficult to identify from morphology alone leading to some mislabelling in living collections.
- Here, we initiate development of a genetic barcode system for the identification of *Fraxinus* species based on low-copy-number protein coding genes. We also conduct experimental crosses among ash species in different sections.
- Three genes are effective in identifying ash samples to sectional level but only in some cases to species level. They highlight that *Fraxinus mandshurica*, *Fraxinus platypoda* and *Fraxinus chiisanensis* may be frequently mistaken for one another in living collections. We succeeded in generating 10 wide hybrid plants: two of section *Melioides* (species: *Fraxinus pennsylvanica*) × section *Fraxinus* (species: *Fraxinus excelsior*) and eight of section *Ornus* (species unclear) × section *Fraxinus*

Disclaimer: The New Phytologist Foundation remains neutral with regard to jurisdictional claims in maps and in any institutional affiliations.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2025 The Author(s). *Plants, People, Planet* published by John Wiley & Sons Ltd on behalf of New Phytologist Foundation.

(species: *F. excelsior*). One hybrid from each of our crosses has survived natural infection with the ash dieback pathogen in Ireland. We also discovered a hybrid between section *Melioides* (species: *F. latifolia*) × section *Fraxinus* (species: *F. excelsior*) formed spontaneously in the ash collection at Kew.

- Our findings facilitate the utilisation of global ash species diversity in response to novel threats and highlight the difficulty of designing a barcoding system capable of distinguishing all species of a genus.

1 | INTRODUCTION

In the temperate regions of Europe, European ash (*Fraxinus excelsior*) is an ecologically and economically important broadleaf tree species (Dobrowolska et al., 2011; Mitchell et al., 2014) but is severely affected by the rapid spread of the non-native fungal species, *Hymenoscyphus fraxineus*, causing symptoms commonly referred to as ‘ash dieback’ (ADB) or ‘Chalara’ (Coker et al., 2019; Gross et al., 2014; Pautasso et al., 2013). European ash populations are further threatened by an insect species, the emerald ash borer (EAB) (Orlova-Bienkowska & Bienkowski, 2018), which has already devastated ash populations in North America (Herms & McCullough, 2014). Worldwide species of ash differ in their levels of resistance to ADB and EAB (Kelly et al., 2020; Kelly et al., 2025; Koch, 2025; Koch et al., 2007; Kowalski et al., 2015; Nielsen et al., 2016; Plumb et al., 2019; Rigsby et al., 2016). Thus, some non-native *Fraxinus* species could be suitable to replace susceptible populations in Europe and North America (Jepson & Arakelyan, 2017; Lévesque et al., 2023; Marzano et al., 2019) or used as parents in hybrid breeding programmes to transfer resistance among species (Koch et al., 2007, 2012, Plumb et al., 2019).

Several classifications of the genus *Fraxinus* have been published (e.g. Hinsinger et al., 2013; Jeandroz et al., 1997; Lingelsheim, 1907; Miller, 1955; Nikolaev, 1981; Wallander, 2008, 2012; Wenzig, 1883; Wesmael, 1892) with differing species divisions and sectional divisions. In this paper, we largely follow the classification of Wallander (2012), where the genus contains over 40 species, many of which are placed into six sections. Our only exception to this is that we do not place *Fraxinus griffithii* into section *Ornus* because of its placement outside the *Ornus* clade in the phylogeny of Kelly et al. (2020) and Hinsinger et al. (2013); we treat it as *Incertae sedis*. We note that Hinsinger et al. (2013) place *Fraxinus platypoda*, *Fraxinus chiisanensis* and *Fraxinus cuspidata* in section *Melioides* s. l.. By contrast, in Jeandroz et al. (1997), *F. platypoda* is placed as a sister to *Fraxinus mandshurica* and *F. cuspidata* is in a polytomy with clades containing species of Wallander's sections *Fraxinus* and *Melioides*. In Kim et al. (2022), *F. chiisanensis* and samples of *F. platypoda* from Japan are placed in *Melioides* s.l., but samples of *F. platypoda* from China are placed in section *Fraxinus* where they are in a clade with *F. mandshurica*. Given these conflicts and uncertainties, we treat *F. platypoda*, *F. chiisanensis* and *F. cuspidata* as *Incertae sedis* (as in Wallander, 2012).

Fraxinus species can be difficult to tell apart as many of their most important morphological features are only visible at certain times of

year or are age dependent (Henry, 1914; Wallander, 2012). Some closely related species are morphologically similar and have been defined partly on the basis of their geographic distribution. Some species show high intraspecific variability and phenotypic plasticity. It is hard to differentiate between some *Fraxinus* species using DNA sequencing of the internal transcribed spacer (ITS) of nuclear ribosomal DNA, as sequence similarity is high between closely related species and intragenomic variation is present within some individuals (Wallander, 2008). Ploidy level (and hence C-value) is a useful method to distinguish among certain species but most species are diploid and others, such as *F. caroliniana*, include more than one ploidy level (Whittemore et al., 2018).

Hybridisation occurs between some species of *Fraxinus*. The sister species *F. excelsior* and *Fraxinus angustifolia* naturally hybridise in Europe (Fernández-Manjarrés et al., 2006; Heuertz et al., 2006; Thomasset et al., 2011) and have also been crossed by breeders (Raquin et al., 2002). Many of the *Fraxinus* species in the section *Melioides* clade hybridise in areas of coexistence in North America (Russell M. Burns, 1990; Wallander, 2008). There is anecdotal evidence of natural hybridisation in regions of species overlap between section *Melioides* and section *Ornus* (Wallander, 2012). Henry (1914) documented various artificial interspecific crosses of *Fraxinus* species, although the extent to which they were successful is unclear. Hybrids of *F. nigra* × *F. mandshurica* (both within the section *Fraxinus*) were produced by Davidson and two commercial clones, *Northern treasure* and *Northern gem*, were patented (Davidson, 1999; Davidson & Ronald, 2001). Koch et al. have successfully produced new *F. nigra* × *F. mandshurica* hybrids and second-generation backcrosses appear to be resistant to EAB (Koch et al., 2007, 2010). Within the section *Fraxinus*, hybrids between *F. mandshurica* and *F. angustifolia* subsp. *syriaca* (syn. *F. sogdiana* used by authors of the study) have been produced and propagated using embryo culture, with one clone showing photosynthetic heterosis (He et al., 2021, 2023). Wright (1953) produced hybrids between *Fraxinus pennsylvanica* and *Fraxinus velutina* (both in section *Melioides*). A cross between *F. velutina* and *F. pennsylvanica* has been used to map candidate loci for salt tolerance (Liu et al., 2024). Wider hybrids between sections *Fraxinus* and *Melioides* have also been produced. Johnson and Heimburger (1946) reported hybrids between *F. excelsior* var. *aureovariegata* (section *Fraxinus*) and *Fraxinus americana* (section *Melioides*); *F. pennsylvanica* (section *Melioides*); and *Fraxinus quadrangulata* (section *Dipetala*). Hybrids have been produced in China involving

F. mandshurica and *F. americana*, and *F. mandshurica* and *F. velutina*, using electrostatic treatments to overcome fertility barriers; these hybrids showed heterosis in growth (Cao et al., 2023; He et al., 2019; Zeng et al., 2015). Hybrids between *F. mandshurica* (section *Fraxinus*) and *Fraxinus chinensis* (section *Ornus*) have recently been produced (Liu et al., 2025).

Hybrid breeding programmes have been set up in response to other well-established pathogens of trees, such as Dutch elm disease (Santini et al., 2007; Smalley & Guries, 1993) and chestnut blight (Anagnostakis, 2012), producing trees with increased resistance. The process of interspecific hybridisation is usually not as simple as identifying two trees of different species and cross-pollinating them. Prezygotic incompatibilities include flowering time differences between the species, failure of pollen germination on the stigma of a different species, and reduction or halting of pollen tube growth. Post-zygotically, the hybrid endosperm or embryos may fail to develop. Interventions may allow some of these barriers to be overcome, including pollen storage, mentor pollen (Knox et al., 1987) and embryo rescue (Sharma et al., 1996; Raquin et al., 2002).

While considerable effort may be invested in producing novel hybrids among tree species, hybrids may form spontaneously in arboreta and botanic gardens (M Maunder, C Hughes, JA Hawkins, A Culham, 2004). Such hybrids have often been viewed as something to be avoided (Ensslin & Godefroid, 2019), especially if they are hard to identify. However, if we can easily discover hybrids and identify their parental species using molecular markers, such spontaneous hybridisation events could become a valuable resource in plant breeding.

The growing interest in research programmes that involve different species of ash and their hybrids (Cao et al., 2023; He et al., 2021; Kelly et al., 2020; Koch et al., 2007, 2012; Liu et al., 2024; Liu et al., 2025; Plumb et al., 2019; Zeng et al., 2015) means that a method for their easy identification, usable on any tissue type at any time of the year, is highly desirable. Whole-genome sequences for 22 *Fraxinus* species have recently been published (Kelly et al., 2020). This provided us with the opportunity to undertake genome-wide searches for nuclear loci that might be suitable for barcoding within the genus. Here, we describe the development of a preliminary set of barcoding regions using single-copy genes in *Fraxinus*. We show that these barcodes can identify specimens to sectional level and often to species level. We also produce new inter-sectional *Fraxinus* hybrids and verify their parentage using our new barcodes.

2 | MATERIALS AND METHODS

2.1 | *Fraxinus* accessions

We collected materials from *Fraxinus* accessions in arboreta and botanic gardens (Table S1) and from the United States Department for Agriculture Forest Service ash breeding programme at the Northern Research Station (in Delaware, Ohio). We focussed on these, rather than herbarium specimens, as they are potential sources for

experiments, future planting stocks and hybridisation programmes. Some of these specimens were used in separate studies as sources of scion material to set up assays for EAB susceptibility in Ohio (Kelly et al., 2020) and trials for ADB susceptibility in England (see: www.forestresearch.gov.uk/research/testing-a-range-of-ash-species-for-tolerance-to-ash-dieback/). The living collections in the British Isles that contributed materials were the following: Royal Botanic Gardens Kew (including Wakehurst), Cambridge University Botanic Garden, Westonbirt Arboretum, Earth Trust, RHS Wisley, Ness Botanic Gardens, Royal Botanic Garden Edinburgh, Dawyck Botanic Garden, The National Botanic Gardens Glasnevin (Ireland), The John F. Kennedy Arboretum (Ireland) and a private collection belonging to Gerry Douglas (Dublin, Ireland). We purchased seeds from Sandeman Seeds (Dalkey, Ireland) labelled as *F. mandshurica* and germinated them in Ireland. We did not have access to some accepted taxa of *Fraxinus*, mainly because of poor representation of species native to Mexico and South America in British and Irish collections. As these species do not grow well in European conditions, they are unlikely to be part of the solution to ADB and EAB. DNA was extracted from the leaves or cambial tissue of specimens using Qiagen DNeasy® Plant Mini Kit (QIAGEN Ltd., Manchester, UK) or CTAB protocols (Kelly et al., 2020).

2.2 | Selection of candidate barcoding genes

A genome assembly for *F. excelsior* was published in 2017 (Sollars et al., 2017) based on a tree grown from seed collected in a woodland in Gloucestershire, UK. For a further 28 of the samples that we used, Kelly et al. (2020) had previously sequenced whole genomic DNA (see Column E in Table S1) and produced genome assemblies. The species identifications of these samples had been checked by Dr Eva Wallander (an expert on the genus) based on morphology and ITS, and where these differed from the labels in the living collections, we took the identification by Wallander to be correct. In the case of accessions 1973–6204 and chi-12, there was uncertainty about the identity and Kelly et al. (2020) initially labelled them as *Fraxinus* sp. From the whole genome assemblies of these 29 *Fraxinus* samples, Kelly et al. (2020) had identified groups of putatively orthologous protein coding gene sequences in them and three outgroups (*Olea europaea*, *Erythranthe guttata* and *Solanum lycopersicum*) using OMA (Altenhoff et al., 2019). Full details of the methods can be found in Kelly et al. (2020), but briefly, the full-length sequences (i.e. including both exons and introns, where present) for OMA groups that included sequences from all 29 *Fraxinus* samples and the three outgroups were aligned using MUSCLE (Edgar, 2004) via GUIDANCE (Sela et al., 2015) and unreliably aligned positions removed. OMA groups with alignments shorter than 300 characters in length, or which included sequences with <10% non-gap characters, were excluded, leaving a set of 1385 OMA groups.

The alignments for these groups were examined using a pre-release version of CONTEXT, a phylogenomic dataset browser (v0.8. pre-release) (<https://github.com/lonelyjoeparker/qmul-genome->

convergence-pipeline/blob/master/CONTEXT.md) to generate statistical information about each OMA group including longest sequence length, longest un-gapped region, number of variant nucleotides and amino acids, and the mean entropy values for nucleotides and amino acids. We selected candidates for further analysis based on their degree of variance (25% or above), greatest overall length and length of the longest un-gapped region (being longer than 800 bp). For these candidate OMA groups, maximum likelihood trees were generated using RAXML-ng version 0.8.0 (Kozlov et al., 2019), using the following parameter settings: --model GTR + G --tree pars{10},rand{10} --bs-trees 1000. We examined the trees for congruence with the species tree from Kelly et al. (2020). We checked for putative paralogues by searching the sequence from *F. excelsior* against its reference genome (assembly version BATG0.5) using BLAST; OMA groups were excluded for if the sequence matched another region of the genome with total coverage of greater than 75% of the query sequence length (suggesting that one or more paralogues of the query gene was present in the genome). The sequences for the three outgroups were removed from the remaining candidate OMA groups and new alignments generated using the MUSCLE tool from EMBL-EBI (<https://www.ebi.ac.uk/Tools/msa/muscle/> accessed October 2017).

2.3 | Primer design and testing

We selected four variable OMA groups for primer design. These were OG8679 (a pentatricopeptide repeat, according to the functional annotation of BATG0.5), OG9972 (a histidine phosphotransferase protein), OG22833 (a protein of unknown function) and OG8143 (a polyadenylate-binding protein 1). Primers were designed using Primer3web v4.0.0 (<http://bioinfo.ut.ee/primer3/> accessed September 2017) and purchased from Eurofins MWG Operon (Ebersberg, Germany). The primers were provided and lyophilised, and Milli-Q® H₂O was used to reconstitute them to a concentration of 100 pmol/μl. We tested these by performing polymerase chain reaction (PCR) on our DNA extractions from 151 trees in living collections.

Each PCR reaction contained <100/ng genomic DNA, 1 μl forward primer 10 pmol/μl, 1 μl reverse primer 10 pmol/μl, 25 μl DreamTaq Green PCR Master Mix (2X) (DreamTaq DNA Polymerase), 2 × DreamTaq Green buffer, dNTPs, 4 mM MgCl₂ (Thermo Fisher) and 10 μl 5 × TBT-PAR (containing trehalose, bovine serum albumin [BSA], polysorbate-20 [Tween-20] and 2 μl DMSO [dimethyl sulfoxide]). Reactions were made up to 50 μl with Milli-Q® H₂O. The conditions used were an initial denaturation stage at 94°C for 6 min, followed by 38 cycles of denaturation at 94°C for 1 min, annealing at 52°C for 1 min and extension at 72°C for 2 min, with a final extension at 72°C for 7 min. PCRs were carried out on a GeneAmp® PCR System 9700 thermocycler (Applied Biosystems, Warrington, UK).

Agarose gel electrophoresis was used to determine the presence of correctly sized PCR products.

For sequencing, the PCR products were purified using a NucleoS-pion® Gel and PCR Clean-up kit (Macherey-Nagel). Sequencing reactions were made up of 20–25/ng of the purified PCR product, 0.5 μl at 5 pmol/μl of appropriate primer, 1 μl 5× sequencing buffer, 0.25 μl

DMSO and 0.25 μl BigDye® Direct Sanger Sequencing Kit (Thermo Fisher Scientific UK Ltd [Loughborough]). Cycle sequencing was performed using the following parameters: 35 cycles of denaturation at 96°C for 10 s, annealing at 52°C for 5 s and extension at 60°C for 4 min using a GeneAmp® PCR System 9700 thermocycler (Applied Biosystems, Warrington, UK). Samples were Sanger-sequenced using a 3730 Series Genetic Analyzer (Thermo Fisher Scientific UK Ltd [Loughborough]). The electropherograms were trimmed, and the forward and reverse sequence for each sample were assembled into contigs using Geneious® ver8.1.9 software (Biomatters, Ltd., Auckland, New Zealand).

We aligned sequence data for the four barcode regions from all successfully sequenced samples using MUSCLE (Edgar, 2004) within JALVIEW (Procter et al., 2021). We trimmed the alignment to the ends of the outermost PCR primers. Each sample was given a label that began with an abbreviation of its taxonomic section (DIP = *Dipetala*, FRA = *Fraxinus*, MEL = *Melioides*, ORN = *Ornus*, PAU = *Pauciflorae*, SCI = *Sciadanthus*, INC = *Incertae sedis*), followed by its species ID and its sequencing tube number. For the samples that had been used for whole genome sequencing and barcode primer design, the label contained the section and species allocated after identification by Eva Wallander using morphology and ITS data. For the other samples, the label contained section and species according to labels in living collections. We manually sorted the alignment by section and species. When this was carried out, it was apparent that in all barcodes there were single nucleotide variant (SNV) alleles specific to sections and some species, but that some samples were anomalous. The anomalous samples were given the prefix ANOM and were reassigned to different sections according to the alleles they contained, with their new section abbreviation placed at the start of their label. Within sections, we then sought to manually cluster species, where possible, according to species-specific SNV alleles. Where a species included no samples that had been whole genome sequenced, we assumed that the majority of living collection labels that matched the correct section were also correct at a species level, allowing us to provisionally assign some of the anomalous samples to species. For each sample, we noted the SNV loci in each barcode that allowed assignment to section and, where relevant, species. Once we had as far as possible identified each sample (that had not been included in the whole genome sequencing panel) using barcodes, we compared these identifications, where possible, to independent identifications of the samples made by Eva Wallander using morphology and ITS, and also flow cytometry assessment of genome size.

2.4 | New hybrid crosses

In order to produce new interspecies crosses, we took scion material from a collection of mature *F. excelsior* plus trees selected from Irish forests for good forestry traits. These were five predominantly female trees (Fex 32, 076, 8X and G 40) and five predominantly males (Fex 359, 42, 405-14 and G1). We took scion material from the John F. Kennedy Arboretum, Co. Wexford, Ireland, from trees labelled as

F. pennsylvanica (JFK.02931), *F. paxiana* (JFK.02928), *Fraxinus floribunda* (JFK.02916), *Fraxinus japonica* (a synonym for *F. chinensis* in Wallander, 2012)(JFK.02920), *F. chinensis* (JFK.02906), *F. chinensis* subsp. *rhynchophylla* (JFK.02908), *Fraxinus retusa henryana* (a synonym for *F. floribunda* in Wallander, 2012)(JFK2005.0132), *F. nigra* Marsh. × *F. mandshurica* (Canadian Hybrid No 8921 'Northern Treasure' JFK.2001.0159) and *F. americana* (JFK.02899). From the National Botanical Gardens of Ireland, Glasnevin, we took scions from: *F. texensis* (a synonym for *F. albicans* in Wallander (2012), XX.011046.A2 and *F. mandshurica* (1934.011053.A2). We also took scion material from trees labelled as *F. mandshurica* (T1-T3) in Dr Gerry Douglas' private collection, which were originally sourced from National Institute of Forest Science in Seoul, South Korea.

Scions were cut to length 15–20 cm and ~0.5 cm diameter, with at least two sets of lateral buds and one terminal bud. These scions were cleft-grafted to rootstocks at Teagasc, Kinsealy, Ireland between 2011 and 2015 as follows. Two year old trees from naturally occurring Irish *F. excelsior* populations were sourced from a commercial nursery. During the period February–March, as the scion and rootstocks lay dormant, the rootstock trees were cut to 5–10 cm above soil level and an incision made to receive the scion. The basal end of the scion was cut to two opposing smooth-tapered cuts approximately 2–4 cm in length from the lowest set of buds using a sharp grafting knife. The cut scions were inserted into the rootstock cleft, with the cambium of the scions positioned into direct contact with the cambium of the rootstock. The graft union was bound using elastic grafting bands. The entire scion and graft union was dipped into molten paraffin wax (45°C). Grafted plants were potted into 3 L pots with commercial-grade peat compost and maintained in the greenhouse. Sprouts that arose from below the graft union were routinely removed as these can lead to graft rejection. Grafted trees were later moved to the Teagasc Ashtown Research Centre, Ireland in 2018, following the Kinsealy site's closure.

Cross-pollination was carried out between March and April of 2015 as some of the grafted trees came into flower. The grafted plants were at least 2 years old at the time of flowering and crossing. The following non-native accessions were used in the crosses: trees labelled as *F. mandshurica* in Dr Gerry Douglas' collection were used as male parents (no female flowers were available), *F. pennsylvanica* (JFK.02931) used as a female parent, *F. chinensis* (JFK.02906) was used as a male and a female parent, *F. retusa henryana* (JFK2005.0132) was used as a female parent, *F. japonica* (JFK.02920) was used as a male parent and *F. nigra* Marsh. × *F. mandshurica* (Canadian Hybrid No 8921 'Northern Treasure' JFK.2001.0159) was used as a female parent. Each cross was with *F. excelsior* as the other parent. In addition to the grafted *F. excelsior* plus trees 076, 8X, G40, 32, 42, 405-14, G1 and 359, we also used *F. excelsior* 98, a selected plus tree which was growing outside nearby.

Pollen was collected from male flowers at anthesis by gently tapping inflorescences so that pollen fell into glass petri dishes (60 × 17 mm; Thermo Fisher). Dishes were labelled with the pollen donors' accession number. Glass was used instead of plastic petri dishes for collection and storage to avoid static electricity making the pollen challenging to manipulate. Pollen was stored for future use by

wrapping parafilm around the petri dish's lid, leaving a small opening for drying. Wrapped dishes were then placed in a glass desiccator containing silica gel and stored in a refrigerator at 4°C until required in the same season. In some cases, when the non-native species were hermaphrodite, the male flowers were emasculated to prevent self-fertilisation. Pollinations were performed at least twice for each cross using pollen collected on the same day or from refrigerator-stored pollen and within a few days of the initial pollination event. Stigmas on selected female trees were checked daily over the period in which the inflorescence was maturing and pollinated when some of them showed a glistening surface. Pollen was applied using a tiny paintbrush that was lightly brushed against the stigmas to release a puff of pollen. A tag was placed around each inflorescence of the pollinated tree, with an allocated number corresponding to the parental details for the cross. In one case, two different pollen donors *F. excelsior* 405-14 and *F. excelsior* G1 were used. Pollen of *F. excelsior* 405-14 was used on the first day and pollen of *F. excelsior* G1 was used on a subsequent day for the same cross as 'mentor pollen' may facilitate hybridisation in some cases. The pollination period was generally in the last 2 weeks of March when stigmas were deemed to be most receptive. After the first pollination event, the trees were moved to a separate section of the greenhouse to minimise potential contamination by nearby ash trees which may have been flowering. After that, the pollinated trees were grown in the glasshouse for the summer period in which unpollinated pistils abscised naturally. As *F. excelsior* plus tree 98 was field grown, the pollinated female inflorescences were enclosed and sealed in pollination bags and unbagged inflorescences were used as open pollinated (OP) intra-species controls under the assumption that any ambient pollen would be from *F. excelsior*.

The pollinated inflorescences were observed for the development of the samaras over several months during which many immature/unfertilised samaras abscised naturally. During the last week of August and the first two in September, an assessment was made of seed production from crosses. Full samaras were excised, and surface sterilised in 7% (w/v) calcium hypochlorite Ca (ClO)₂ solution for 20 min, followed by three rinses in sterile water and aseptic air drying. Thereafter, the samaras were dissected aseptically under a stereomicroscope, and the seeds were removed and distributed onto B5 medium for germination (Gamborg et al. 1968) in small Petri plates (5 cm) containing 6–7 ml of medium with sucrose in concentrations of 2%, 3%, 4% and 5% (w/v), and solidified with phytigel 3 g l⁻¹. Culture conditions were at 22 ± 1°C with 16-h photoperiod using Philips® cool white fluorescent tubes giving a photon flux density of 58.6 μmol/m/s. Seed germination and viability were assessed after approximately 6 weeks, and after this period, ungerminated seeds were either transferred to fresh media, or the embryos were excised from seeds for embryo rescue culturing. In some cases, we did not attempt whole seed germination but embryos were excised and cultured directly after seed collection and surface sterilisation. Embryo rescue was performed under a stereomicroscope in a laminar flow hood under aseptic conditions. An incision was made along the longitudinal axis of the seed and into the endosperm and the outer seed coat was peeled off. The seed was prised open at the

cotyledon end to reveal the embryo attached within one half of the seed. The embryo was gently prised off the endosperm using the tip of a scalpel and placed directly into B5 medium. Whole seeds and isolated embryos were transferred to fresh media every 6–8 weeks at least six times. We routinely used B5 medium with 3% sucrose in the latter culture periods. Germinated seedlings were transferred to glass jars (volume, 150 ml) on WPM medium salts (Lloyd and McCown 1980), and vitamins of B5 medium (Gamborg et al. 1968) supplemented with 3% w/v activated charcoal and then weaned to the glasshouse.

In 2018, leaf material was collected from each hybrid for DNA extraction. The *Fraxinus* genetic barcodes were used to identify successful new hybridisations. DNA extraction, PCR and sequencing were performed in the same way as they were for the living collections (see above discussion). Sanger sequence reads were then trimmed, cleaned and assembled into contigs. The sequence chromatograms were viewed using Geneious prime version 8.1, and we looked for double peaks (i.e. heterozygosity) at single nucleotide polymorphism (SNP) loci that differed between the putative parental species sequenced in the test panel. We carried out an initial screening using the barcode region OG9927, and hybrids found with this locus were also sequenced with the barcoding regions OG22833 and OG8143. From 2018 onwards, the new hybrids were grown in an area of high ADB disease pressure at the Teagasc Ashtown Research Centre, Ireland, and records were kept of death of trees because of ADB.

2.5 | Seeking spontaneous hybrids

As hybrids may spontaneously form in living collections, we collected four volunteer seedlings growing within the main ash collection area at Kew Gardens and used the barcodes to test if they were hybrids.

2.6 | Ploidy investigations

Where polyploidy has been reported for certain *Fraxinus* taxa, measurement of genome sizes using flow cytometry can aid in the identification of species and their hybrids. In such cases, nuclear DNA contents were estimated following the one-step flow cytometry procedure (Dolezel et al., 2007). Approximately 1 cm² of the sample leaf material was incubated for 30 s on ice in 1 ml of 'general-purpose buffer' (GPB) (Loureiro et al., 2007) supplemented with 3% PVP-40, after which leaf material from the calibration standard *Petroselinum crispum* or *Pisum sativum* was added (Obermayer et al., 2002). This material was chopped together rapidly using a fresh razor blade. Another 1 ml of the isolation buffer was then added to the chopped material, and the homogenate was filtered through a 30- μ m nylon mesh (CellTrics 30 μ m mesh, Sysmex, Goritz, Germany). Following filtering, 100 μ l propidium iodide (1 mg/ml) was added, and the sample was incubated on ice for 10 min before analysis. The relative fluorescence of 5000 particles was then recorded using a Partec Cyflow SL3 flow cytometer (Partec GmbH, Münster, Germany) fitted with a 100 mW green solid-state laser (532 nm, Cobolt Samba, Solna, Sweden), and the output histograms were analysed with the FlowMax software v.2.4 (Partec GmbH, Münster, Germany).

3 | RESULTS

3.1 | Testing of barcodes

At least one barcode region (Table 1) amplified successfully and yielded sequence data for 148 DNA samples from 45 *Fraxinus* species or subspecies (see Table S2). Usable sequences were generated for 143 samples with OG9927, 135 samples with OG22833, 133 samples

TABLE 1 Primers designed and tested for *Fraxinus* species barcoding.

<i>Fraxinus excelsior</i> position and gene name	Primer name	Primer sequence (5'-3')	Tm ^a	Expected amplicon length
Contig238 91392-93122 Histidine phosphotransferase protein OG9927	9927(1)fwd	TGGAAGGCAGAGTTTCTAGGG	59.8°C	~800 bp
	9927(1)rev	GAGCAGAGTGMTGCATTGAA	56.3°C	
	9927(2)fwd	GCTTGGGCRTTGTCAAAGAT	56.3°C	~800 bp
	9927(2)rev	TTAGCATRGTATGCCCCAGT	56.3°C	
Contig5804 11231-10691 unknown function OG22833	22833fwd	TTGACATGGGATTYAGATGT	54.9°C	~500 bp
	22833rev	ACCATAAAGCTGCACAACCA	55.9°C	
Contig803 88622-87665 Polyadenylate-binding protein 1 OG8143	8143fwd	CACCTCGAACCTTAYCTGCT	58.3°C	~800 bp
	8143rev	TCTCGYAGTCTTCRATGGT	57.3°C	
Contig382 73390-74770 Pentatricopeptide repeat OG8679	8679(1)fwd	CAAAAACCAAGCSGAAATC	53.2°C	~800 bp
	8679(1)rev	TTCCCATCTTYCTCATGCTC	56.3°C	
	8679(2)fwd	ACRCTTTTGACCATGTATGTTG	55.6°C	~800 bp
	8679(2)rev	GCCTGTCTGAAAACCCAAGT	57.3°C	

Note: Each primer's location is given based on the *F. excelsior* BATG0.5 genome. OG numbers refer to the Ortholog group number assigned to genes in Kelly et al., 2020.

^aTm, melting temperature.

with OG8143 and 132 samples with OG8679. If DNA quality was poor, the second primer sets for OG9927 and OG8679 tended to produce lower-quality reads. Trimmed alignments of all clean sequences generated can be found in Data S1–S4, ordered by section and species. These are also available at [10.5281/zenodo.10066755](https://doi.org/10.5281/zenodo.10066755). Examination of these alignments showed that all the barcodes contained SNV alleles that were specific to taxonomic sections and present in all samples within those sections. There were also many SNV alleles that were specific to species, although these were not always present in all samples from a species.

Table S1 gives a list of all tree accessions used and details of the extent to which they can be identified to section and species levels using each DNA barcode. Barcodes OG9972, OG22833 and OG8143 gave conflicting sectional results for only two samples: these were both cases where two genes agreed and the third gave a result suggestive of a hybrid (Kew 1992-375 and Wakehurst 1994-52). In contrast, barcode OG8679 gave many double peaks in chromatograms and gave sequence results that sometimes conflicted with the other barcodes for nine samples. In the sequence alignment for OG8679, several SNP loci had three or even four alleles present among the samples. To give one example, site 546 was: G in the majority of samples, T in all four samples of *F. paxiana*, one of eight individuals of *F. floribunda*, one of six individuals of *F. excelsior*, C in two of three individuals of *F. lanuginosa*, A in all individuals later identified as *F. chiisanensis* (except one which was R [A/G]) and one of eight individuals of *F. quadrangulata* and in two of four individuals of *F. platypoda*, R in one of seven *F. nigra* and one of five *F. latifolia*. Those individuals with an A at position 546 all had an A at position 592, a G at position 657 and a T at position 674. Those that had an R at position 546 had an A or R at position 592, a K at position 657 and a T or Y at position 674. Such a pattern is best explained by the presence of two paralogues, with some samples showing amplification of both, and some only amplifying one copy, potentially because of loss or mutation of the primer site at the other copy. There were no cases in which OG8679 was able to give us sectional or species assignments for a sample that were unavailable from one of the other barcodes. Therefore, we decided to discard OG8679 as a barcode and focus only on OG9972, OG22833 and OG8143.

Using OG9972, OG22833 and OG8143, 148 samples could be assigned to section (taking a majority vote in the two cases of sectional conflict) and 88 could be assigned to species. Those 60 samples that could not be identified to species level were all in the sections *Melioides*, *Fraxinus*, *Dipetala* or *Ornus*. These sections are known to have similar species that sometimes hybridise. No diagnostic alleles were found for *F. excelsior* (in section *Fraxinus*), the most widespread species in Europe, or *F. pennsylvanica* and *F. americana* (in section *Melioides*), the most widespread species in North America, probably because of widespread hybridisation with other species found within their respective ranges.

Of the 148 samples assigned to section or species level by the three barcodes, 67 were also identified by Eva Wallander using morphology and ITS. Of these, none were assigned to a different section by the two methods of identification. For 34 samples, we gained species level

identifications using both methods. In 38 cases, these agreed. The cases of disagreement were for *F. pennsylvanica* / *F. caroliniana* / *F. velutina* which are all closely related within section *Melioides*, or *F. chinensis* / *F. baroniana* / *F. floribunda* in section *Ornus*.

All three barcodes were each able to identify most of the 148 samples to sectional level. However, section *Fraxinus* was hard to identify using OG22833 and OG8143. For section *Fraxinus*, OG9972 performed best, with three clear section-specific alleles (Table 2). Overall, OG9972 identified 54 samples to species level, OG22833 identified 65 samples to species level and OG8143 identified 47 samples to species level. Within section *Dipetala*, OG9972 performed best at identifying species and OG8143 could not distinguish any species. Within section *Fraxinus*, OG9972 and OG22833 both performed better than OG8143, but OG8143 was particularly useful for distinguishing *F. mandshurica*. In section *Melioides*, all barcodes were unable to distinguish species, with the exception of *F. velutina*, which could be identified by OG22833. In section *Ornus*, the barcodes differed widely in their patterns of diversity: OG9972 was good for identifying *F. floribunda* and *F. lanuginosa*; OG22833 was good for distinguishing *F. chinensis*; and OG8143 was good for distinguishing *F. chinensis* and *Fraxinus ornus*. Within the small section *Pauciflorae*, all barcodes could distinguish *Fraxinus greggii* but only OG8143 could distinguish *Fraxinus gooddingii*. All barcodes performed well at distinguishing the species not assigned to sections (*Incertae sedis*), but OG8143 was alone in being unable to identify *F. griffithii* to species level.

Of our 148 sample identifications with barcodes, 25 conflicted with the labels in the living collections that they were derived from. Nineteen of these conflicts involved assignments of different sections, and six were assignments of different species within sections. Of the 19 sectional conflicts, four identifications were also available from Eva Wallander who gave sectional classifications that agreed with the barcodes.

There were two cases where the tree label said *F. platypoda* (*Incertae sedis*), but the barcodes indicated *F. mandshurica* (*Fraxinus*). These were JFK.02932 and USDA FS plat-3, and for the latter, the barcode identification was confirmed by Wallander. The barcodes assigned Kew 1900-44501 *F. quadrangulata* (*Dipetala*) and Kew 2011-1622 *F. quadrangulata* (*Dipetala*) to section *Fraxinus*. A sample labelled *F. greggii* (Cambridge 19860253*A) was identified as *F. cuspidata* (*Incertae sedis*) by the barcodes. A sample labelled *F. chinensis* subsp. *chinensis* (*Ornus*) (Westonbirt 56.0455) was identified as *F. pennsylvanica* (*Melioides*) by both the barcodes and Wallander. A sample labelled *F. platypoda* (*Incertae sedis*) (USDA FS plat-1) was identified as *F. pennsylvanica* (*Melioides*) by both the barcodes and Wallander (it is now relabelled in the USDA FS collection as pe-103). Three samples labelled as *F. nigra* (*Fraxinus*) (Wakehurst 1996-5167 and 1996-5169 in replicate) were identified as section *Melioides* by the barcodes, and 1996-5169 was identified as *F. pennsylvanica* (*Melioides*) by Wallander. One sample labelled *F. americana* (*Melioides*) (JFK.02899) was identified as *F. chinensis* (*Ornus*) by the barcodes; morphological evaluation of this accession is needed. Seven samples labelled as *F. mandshurica* (GD-T3, GD-T1 and Kew 1989-3691, 1989-8285, 1992-375, 1992-376 and 1992-377) in section *Fraxinus* were identified by the barcodes as being not in the section *Fraxinus*.

TABLE 2 Diagnostic single nucleotide polymorphism (SNP) allele locations for sections and species of the genus *Fraxinus* using three barcodes.

Section	Species	OG9972 SNPs	OG22833 SNPs	OG8143 SNPs
Dipetala		100, 342, 1093, 1405–1407	147, 182, 419	109, 122, (227), 303, 484, 522
	<i>F. anomala</i>	1238		
Fraxinus	<i>F. quadrangulata</i>	(268), (498), (1057)	(443), (465)	(529)
		618 + 743, 1482, (347), (1538)	66–67 + 362–364 + 366–373, (352)	304, (360), (648)
	<i>F. angustifolia</i>	(805), (897), (1256)	(369), (412)	
	<i>F. mandshurica</i>	(306), (452), (574), (749)	324	(129), (239) (262), 315, (767)
	<i>F. nigra</i>	1663	(275), 340, 425, 446	
Melioides		603, 648, 839, 955, 1048, 1149, 1193	144, 180, 366	200, 237, 757
	<i>F. caroliniana</i>	676		
Ornus	<i>F. velutina</i>		24, 110	
		381, (1374)	66, 67, (255)	(156), (174), (205), (249), (321), (374), (676)
	<i>F. chinensis</i>			(156)
	<i>F. floribunda</i>	(207), (361), (1605)	(194)	(427), (750)
	<i>F. ornus</i>			169
Pauciflorae	<i>F. lanuginosa</i> and <i>F. sieboldiana</i>	293, 316, 567, 729, 1071, 1163, 1470, 1701, 1716	275, 364, 426	109, 526, 535
	<i>F. greggii</i>	1686	160, 190, 223	725
Sciadanthus		698, 1577	338, 381, 509	782
	<i>F. xanthoxyloides</i>	(1619)		
Incertae sedis		(153), (241), (463), (603), (624), (764), (1091), (1185), (1187), (1563)	66–67 + 144 + 366 + 372 + 272	(390), 709
	<i>F. chisanensis</i>			
	<i>F. cuspidata</i>	313, 319, 323, (537), 1,152, 1335, 1412	45, 182, 223, 359, 415	86, (173–226), 263, (340), 421, 495, 559, 621
Incertae sedis	<i>F. griffithii</i>	150, 408, 490, 894, 1140–1154, 1185, 1468, 1496, 1564	43, 76, 219, 362, 469, 494	
	<i>F. platypoda</i>	1438, 1469	38	182, 409

Note: Locations refer to positions in the alignments in Data S1–S9, where the first position in the first base in the forward primer. Locations in brackets mean that there is an allele unique to the species but not fixed within the species, so its presence is diagnostic but its absence is not. The plus sign is used for loci that must be in a particular combination to identify a species or section. The sections are according to Wallander (2013) classification with the exception of *F. griffithii* which we place as *Incertae sedis* because of its placement in the species tree inferred in Kelly et al. (2020).

Closer examination of their genome size and bud morphology (see details below) allowed them to be identified as *F. chiisanensis* (*Incertae sedis*). Five seedlings germinated in Ireland from seeds purchased from Sandeman seeds (Dalkey, Ireland) labelled as *F. mandshurica* were confirmed to be *F. mandshurica* by barcodes.

Of the six cases where the barcodes assigned a different species within the same section as the living collection label, three of them were also examined by Wallander who in two cases made the same identification as the label (USDA FS PE_48 FRAX10 *F. pennsylvanica* which the barcodes assigned to *F. velutina* and USDA FS Bar-2 FRAX28 *F. baroniana* which the barcodes assigned to *F. chinensis*). In one case, Wallander made the same identification as the barcodes (Kew 1973-6204 FRAX29 *F. bungeana* which the barcodes assigned to *F. ornus*), suggesting that the barcoding was accurate. The other three cases involved JFK.02906 *F. chinensis* being assigned to *F. ornus*; Kew 2004-1231 *F. paxiana* and USDA FS BUN-5 *F. bungeana* being assigned to *F. chinensis*. Details of the SNP loci used for each assignment can be found in Table S2 and seen in Data S1–S3.

F. excelsior (USDA FS ex-12) had several positions in OG9927 that were diagnostic for *F. cuspidata* (*Incertae sedis*) or were heterozygous for a base found in section *Fraxinus* and a base found in at least one individual of *F. cuspidata*, suggesting it could be a hybrid between these two species. However, in OG22833 and OG8143, there was no evidence of this.

3.2 | Genome size measurements

Genome sizes were measured for 49 *Fraxinus* accessions. These results are shown in Table S1, column T, with inferred ploidy level from these genome sizes in column U. The majority of samples had 1C genome sizes between 700 and 1000 Mb and were inferred to be diploid. Some, but not all, accessions identified by barcodes as *F. chinensis* were hexaploid: Westonbirt 5.0456, USDA FS chi-16, JFK.02920, JFK.02908 and 2GD were hexaploid but JFK.02899 and USDA FS bar-2 (FRAX 28) were diploid. Kew's 1992-377 and the samples in Gerry Douglas' collection, all labelled as *F. mandshurica* but with anomalous barcode sequences, were hexaploid (unlike JFK 02932 identified by barcodes as *F. mandshurica* which had a diploid C-value). The other polyploids found were all samples for which the barcodes had failed to sequence: *Fraxinus biltmoreana* Westonbirt 5.0463 (FRAX17) was hexaploid, *Fraxinus profunda* Wakehurst 1993-1493 appeared to be octoploid, *F. lanuginosa* RBGE 20071296-G (FRAX22) was tetraploid, and *Fraxinus uhdei* USDA FS uhdei-ponto (FRAX34) was hexaploid; it is likely that the barcodes did not work well in these samples because of divergent homologous copies of the barcode sequences among their sub-genomes.

3.3 | *F. chiisanensis*

Closer analysis of the morphology of the adult samples labelled as *F. mandshurica* at Kew and in Gerry Douglas's collection, but with

anomalous barcodes and hexaploidy, showed them to have naked buds which is characteristic of *F. chiisanensis* (Wallander, 2012), a rare endemic of Korea (Kim et al., 2022). The seeds for the adult trees in Gerry Douglas' collection were sourced from the National Institute of Forest Science in Seoul, South Korea. A chromosome squash on a seedling from one of the Kew trees, which was hermaphroditic (1992-377), was also consistent with hexaploidy. A previous unpublished study also suggested that *F. chiisanensis* is hexaploid (mentioned in Siljak-Yakovlev et al., 2014). Interestingly, the pattern of barcode polymorphism within these putative *F. chiisanensis* samples showed some unique alleles (OG9972: 1563, 1091, 1187, 764; OG8143: 390, 709) and also patterns of polymorphism involving alleles common in different sections (OG9972: 123, 153, 381, 405, 1048, 1487, 1698). In OG8143, *F. chiisanensis* shared alleles with *Melioides* at some loci (OG8143: 262-267, 801) but shared alleles with everything but *Melioides* at other loci (OG8143: 200, 237, 736-738). In OG22833, *F. chiisanensis* shared alleles with everything but section *Ornus* at 66-67, everything but *Melioides* at 144, everything but *Melioides* and *Dipetala* at 366, everything but *Ornus* and *Sciadanthus* at 372 and everything but *Pauciflorae* at 272. Thus, within its six genome copies, *F. chiisanensis* harbours a complex mix of variation found scattered across other sections of the genus.

3.4 | Hybrid crosses

Because of the barcoding and flow cytometry results reported earlier, several of the trees that were grafted and used for hybridisation were re-identified, including some that were used as parents in our crosses. *Fraxinus chinensis* (JFK.02906) was identified as diploid *F. ornus*. *F. japonica* (JFK.02920) (which, as noted above, is a synonym for *F. chinensis* (Wallander, 2012)) was identified as the hexaploid *F. chinensis*. The adult trees labelled *F. mandshurica* in Dr Gerry Douglas' collection were identified as hexaploid *F. chiisanensis*. *Fraxinus retusa henryana* (a synonym for *F. floribunda* in Wallander, 2012) (JFK2005.0132) was confirmed as a diploid in section *Ornus* but could not be assigned to a species; we cannot be sure if it is *F. floribunda* or *F. ornus*. *Fraxinus pennsylvanica* (JFK.02931) was confirmed as being in section *Melioides* but could not be assigned to a species by the barcodes; we, therefore, have no reason to believe that the label is incorrect. *Fraxinus nigra* Marsh. × *F. mandshurica* (Canadian Hybrid No 8921 JFK.2001.0159) was not examined using barcodes, but we are confident that its parentage is reliably known (Davidson & Ronald, 2001).

Of the 35 crosses and controls that were performed (Table S2), 756 seeds were produced, of which 363 were propagated, producing 131 seedlings of which 85 survived to be viable plants. We carried out an initial screening using the barcode region OG9927 on 84 seedlings that were alive at the time of the screening. This identified 16 putative hybrids showing heterozygosity at sites that varied among parental species. We then also sequenced these individuals with the barcoding regions OG22833 and OG8143. Region OG2833 was found to produce clear sequence reads of a comparable length

to the average sequence length for the pure *Fraxinus* species. However, for the barcode region OG8143, we found only the forward primer was able to produce useable sequence reads for all of the putative hybrids, and the useable region after sequence clean-up was found to be ~650 bp; this is ~100 bp short of the length of the full region.

None of the crosses that used non-native species as pollen donors produced hybrid offspring. In all of these cases, the offspring were sired by *F. excelsior* pollen. This includes all crosses involving *F. chiisanensis*, and flow cytometry confirmed that the progeny of the *F. excelsior* mother tree used in this cross were all diploid. There were 15 crosses performed with non-native species as the maternal parent. Four of these produced no progeny. Seven of them had progeny that were confirmed as hybrid. We identified 14 intersectional hybrids; this included two plants of *F. pennsylvanica* × *F. excelsior*, six of *F. ornus* × *F. excelsior* and five section *Ornus* × *F. excelsior*. For the latter five, the barcodes did not resolve whether one parent was *F. ornus* or *F. floribunda* within section *Ornus*.

Both of the *F. pennsylvanica* × *F. excelsior* hybrids displayed vigorous growth and had no ash dieback symptoms in 2021, despite growing in an area with prevalent spore inoculum since 2018. One of these was still alive and healthy in July 2023. Most of the hybrids produced between *F. ornus* or section *Ornus*, and *F. excelsior* died of ash dieback symptoms. Four were dead by the end of summer 2020. In 2021, one had severe symptoms, three had mild symptoms, and three were healthy. One was still alive and healthy in July 2023.

Sequence data for these two surviving hybrids for OG22833 and OG9927 are included in Data S1 and S2. For OG9927, in the *F. pennsylvanica* × *F. excelsior* hybrid, positions 416, 454, 603, 648, 743, 839, 1148, 1193 and 1482 were heterozygous. These sites are all diagnostic for section *Fraxinus* or section *Melioides* (see Table 2), confirming that this individual is a hybrid. In the section *Ornus* × *F. excelsior* hybrid, positions 1148, 1465, 1482, 1538 and 1660 are heterozygous, confirming its hybridity between section *Ornus* and section *Fraxinus*. For OG22833, in the *F. pennsylvanica* × *F. excelsior* hybrid, positions 144, 180 and 366 were heterozygous, which are diagnostic for *Melioides* (Table 2) and position 272. In the section *Ornus* × *F. excelsior* hybrid, positions 66 and 67, which are diagnostic for section *Ornus* (Table 2), were heterozygous, as were positions 116, 266 and 332 which are fixed in one of the sections but polymorphic in the other.

3.5 | Spontaneous hybrids

We generated sequence data for four volunteer seedlings collected from the main ash area at Kew. Two of these were identified as *F. ornus* by at least one barcode (Table S2). Another was identified as being from Section *Melioides*. The fourth, which was a germinated seed lying in the grass when first collected, was identified as a hybrid between section *Fraxinus* and section *Melioides* by all three barcode regions and by OG8679. Two barcode regions identified *F. latifolia* as one parent. The seedling was collected close to *F. latifolia* 1981–

8204, so this is likely to be a parent. The other parent was harder to identify, but in barcode OG9927 it contained an allele found only in *F. excelsior* and *F. mandshurica*. In OG8679, it contained an allele found only in *F. excelsior*. Therefore, we conclude that this individual is a hybrid between *F. latifolia* and *F. excelsior* (though we do not know the direction of the cross). It has survived its first summer outdoors in an ash dieback infested area without ash dieback damage but has not yet been fully challenged.

4 | DISCUSSION

We have developed a preliminary set of three low-copy nuclear gene barcodes that can delineate taxonomic sections and in some cases, individual species within the genus *Fraxinus*, allowing identification of ash samples and verification of hybrids. The cases where the barcodes did not perform well were certain polyploid species, and within large sections that contain many closely related species with a shared geographic distribution. For identification of the latter, the use of further low copy number genes, microsatellites (Liu et al., 2025), whole genome sequencing (Kelly et al., 2020; Liu et al., 2024), amplified fragment length polymorphism (AFLPs, Hisinger et al., 2014) or SNP arrays may help. The large amount of gene flow that occurs among some species of *Fraxinus* may still mean that reliably diagnostic alleles are rare. Knowledge of genome-wide variation within many *Fraxinus* species is currently limited, but where it has been surveyed in the case of *F. excelsior*, *F. angustifolia* and *F. mandshurica*, it has shown widespread sharing of alleles (Hisinger et al., 2014). Barcoding of larger sample sizes in the future could identify more alleles found exclusively in a subset of populations within some species.

It is not uncommon for DNA barcodes to be unable to distinguish all species within a plant genus. For six genera within the Asteraceae, Gao et al. (2010) found ITS2 to be the most effective of five tested barcodes with success at species level within each genus varying from 48% to 97%. In the genus *Corydalis*, Ren et al. (2019) gained a maximum of 70% success in species identification, using ITS and matK, and tested three other barcodes that performed less well. In the genus *Curcuma*, the highest rate of successful species identification was 48% by ITS2, out of five barcodes tested (Chen et al., 2015). Bratzel et al. (2020) tested the low-copy nuclear gene *Agt1* as a barcode within the Bromeliaceae and correctly identified 22 of 41 species tested. We publish the loci contained in this paper for the scientific community as a starting point for barcoding in *Fraxinus*, and we hope that improvements will come as more samples are added to the data available, the taxonomy of the genus is further clarified, and further genes are added.

Use of these barcodes suggests that mislabelling is not uncommon in living collections of *Fraxinus*, expanding on earlier observations by Koch et al. (2010) and Koch et al. (2012). Potential mislabelling of Asiatic species may reflect lower acquaintance of European curators with Asiatic species than with European and American species. Several cases of mislabelling in living collections involve a local species being given an exotic label: in some cases, this may be because exotic scion material had been grafted to a local species rootstock, but shoots from

the rootstock have become dominant and the exotic material has not survived. Other cases of mislabelling may be ad hoc errors within collections. Our findings highlight the need for verification of living collection labels before undertaking experiments or initiating breeding programmes. The rare Korean endemic *F. chiisanensis* is commonly labelled as *F. mandshurica*, although a seedlot of *F. mandshurica* supplied by Sandeman Seeds (Dalkey, Ireland) was found to be correctly labelled. Samples we identify as *F. mandshurica* are sometimes labelled as *F. platypoda*, but there seems to be genuine ambiguity around the latter species, as Kim et al. (2022) found Chinese samples of *F. platypoda* to cluster among *F. mandshurica* samples (see also Jeandroz et al., 1997), whereas Japanese samples of *F. platypoda* were in a clade sister to *F. chiisanensis*.

We generated and verified wide hybrids of ash, among sections *Fraxinus*, *Melioides* and *Ornus*: these were from *F. pennsylvanica* × *F. excelsior* and section *Ornus* × *F. excelsior* crosses. We also discovered a spontaneous hybrid formed between *F. latifolia* (section *Melioides*) and *F. excelsior* (section *Fraxinus*) at Kew Gardens. Many hybrids have been reported or made previously within the genus *Fraxinus* (see Section 1). The wide hybridisations recently conducted between sections *Fraxinus* and *Melioides* (*F. mandshurica* × *F. americana*, and *F. mandshurica* × *F. velutina*) Zeng et al., 2015; He et al., 2019), used electrostatic treatments to overcome fertility barriers. We did not use electrostatic treatments to produce the hybrids reported here, nor did Liu et al. (2025) in producing hybrids between *F. mandshurica* (section *Fraxinus*) and *F. chinensis* (section *Ornus*).

One of our major objectives when setting up crosses was to hybridise *F. excelsior* with east Asian ash species possessing high resistance to both EAB and ADB. We have not achieved this, partly because of initial misidentification of parental trees; some of which were labelled as Asiatic species but turned out not to be. *Fraxinus chiisanensis* was believed to be *F. mandshurica* at the time of crossings. We attempted many crosses with *F. chiisanensis* pollen, but these were unsuccessful. We might have had more success by using *F. excelsior* as a pollen donor, because hybridisation and introgression can be asymmetric, as in oaks (Petit et al., 2004); however, no female trees of *F. chiisanensis* were available. We cannot exclude the possibility that there are intrinsic incompatibilities between *F. excelsior* and *F. chiisanensis*. While *F. excelsior* is a diploid, *F. chiisanensis* is hexaploid, and our barcodes suggest it could have a complex hybrid origin.

While the majority of our hybrid crosses died from ash dieback between 2018 and 2023, one individual from each of our crosses remained alive and healthy in summer 2024. Although *F. pennsylvanica* is highly susceptible to EAB (Koch et al., 2015; Orlova-Bienkowskaja & Bienkowski, 2016), it does appear to be less susceptible to ADB than *F. excelsior* Nielsen et al., 2016; Drenkhan et al., 2017; Plumb et al., 2019. Thus, crosses between *F. pennsylvanica* and *F. excelsior* may have promise for ADB resistance breeding, especially as *F. pennsylvanica* is a good timber species. Section *Ornus* contains species with low susceptibility to ADB and EAB Nielsen et al., 2016; Drenkhan et al., 2017; Plumb et al., 2019; Kelly et al., 2020), so hybrids from this section may also be of interest. The spontaneous hybrid between *F. latifolia* and *F. excelsior* has

survived its first summer outdoors in an ash dieback infested area without incurring ash dieback damage. Although *F. latifolia* is highly susceptible to EAB (Kelly et al., 2020), it appears to be less susceptible to ADB than *F. excelsior* (Nielsen et al., 2016; Drenkhan et al., 2017; Plumb et al., 2019), so the hybrid may have some promise. If ADB resistant hybrids are discovered, they may have potential as bridges for gene exchanges between species of interest followed by backcrossing to the main species of interest. Further testing would also be needed, such as to assess their utility for wood production and provision of ecosystem services. Their potential to become invasive or damage the native gene pool would also require careful monitoring.

Spontaneous hybrids in living collections are often seen as a liability and a concern for ex situ conservation (Ensslin & Godefroid, 2019; Maunder et al., 2004). However, with means to readily identify them and their parentage, they could be a valuable resource. Their discovery may save much effort in setting up artificial pollinations and breaking seed dormancy. Here, in a sample of just four volunteer seedlings from Kew Gardens, we identified an intersectional hybrid between the American species *F. latifolia* and the European species *F. excelsior*. Wider sampling of seeds and seedlings in botanic gardens and arboreta may yield many more hybrids.

The barcodes and hybrids presented here provide preliminary methodological and biological resources for the use of world-wide ash germplasm for resistance to pests and pathogens and adaptation to climate change. The success of wide hybridisations suggests that mobilisation of genetic diversity throughout the genus in hybrid breeding programmes may be possible for the genus *Fraxinus*. The development of hybrid breeding programmes between *F. excelsior* and non-native species may be part of the solution to the threats of ADB and EAB. In addition, the more extensive field testing of confirmed ash species (using barcodes) from diverse regions of provenance may identify adapted material for commercial timber production and rapid carbon sequestration.

AUTHOR CONTRIBUTIONS

Richard J. A. Buggs, William J. Plumb, Laura J. Kelly, Jennifer Koch and Gerry C. Douglas planned and designed the research. William J. Plumb, Joe Mullender, David Carey, Laszlo Csiba and William Crowther extracted the DNA. William J. Plumb and Laszlo Csiba conducted Sanger sequencing. Robyn F. Powell conducted flow cytometry. Gerry C. Douglas conducted hybridisations. William J. Plumb, Laura J. Kelly, Miguel Nemesio-Gorriiz, Mary E. Mason and Jennifer Koch collated samples. Richard J. A. Buggs, William J. Plumb, Laura J. Kelly, Jennifer Koch and Gerry C. Douglas wrote the manuscript.

ACKNOWLEDGEMENTS

We thank Eva Wallander for her expert identifications of numerous samples used in this study. We thank Jaume Pellicer for advice and assistance with flow cytometry. We thank Timothy Baxter, Samuel Brockington, Peter Brownless, Dan Crowley, Dawn Edwards, Simon Honey, Ross Irvine, Richard Jinks, Penny Jones, Simon Toomer, Tony Kirkham, Hugh McAllister, Iain Parkinson and Sara Redstone for their help with obtaining *Fraxinus* materials from UK collections. We thank

Colin Kelleher, National Botanic Gardens of Ireland, Glasnevin for provision of scion wood of exotic species and John McNamara for technical assistance in grafting. R.J.A.B. and L.J.K. acknowledge funding from the Erica Waltraud Albrecht Endowment Fund, Department for Environment, Food and Rural Affairs (Defra) Future Proofing Plant Health, and the Living with Environmental Change Tree Health and Plant Biosecurity Initiative – Phase 2 (grant no. BB/L012162/1), funded jointly by BBSRC, Defra, ESRC, Forestry Commission, NERC and the Scottish Government. W.J.P. was supported by a Walsh Fellowship 2014001 from Teagasc and funding from Defra.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are openly available in Zenodo at [10.5281/zenodo.10066755](https://doi.org/10.5281/zenodo.10066755).

ORCID

William J. Plumb  <https://orcid.org/0000-0002-1498-0113>

Laura J. Kelly  <https://orcid.org/0000-0003-1159-939X>

Laszlo Csiba  <https://orcid.org/0009-0008-4520-5807>

Miguel Nemesio-Gorrioz  <https://orcid.org/0000-0001-8052-8101>

David Carey  <https://orcid.org/0000-0001-5616-4379>

Mary E. Mason  <https://orcid.org/0000-0001-9566-4798>

Jennifer Koch  <https://orcid.org/0000-0002-4281-9669>

Richard J. A. Buggs  <https://orcid.org/0000-0003-4495-3738>

REFERENCES

- Altenhoff, A. M., Levy, J., Zarowiecki, M., Tomiczek, B., Warwick Vesztrocy, A., Dalquen, D. A., Müller, S., Telford, M. J., Glover, N. M., Dylus, D., & Levy, J. (2019). OMA standalone: Orthology inference among public and custom genomes and transcriptomes. *Genome Research*, 29, 1152–1163. <https://doi.org/10.1101/gr.243212.118>
- Anagnostakis, S. L. (2012). Chestnut breeding in the United States for disease and insect resistance. *Plant Disease*, 96, 1392–1403. <https://doi.org/10.1094/PDIS-04-12-0350-FE>
- Bratzel, F., Heller, S., Cyrannek, N., Paule, J., Leme, E. M. C., Loreth, A., Nowotny, A., Kiefer, M., Till, W., Barfuss, M. H. J., Lexer, C., Koch, M. A., & Zizka, G. (2020). The low-copy nuclear gene *Agt1* as a novel DNA barcoding marker for Bromeliaceae. *BMC Plant Biology*, 20, 111. <https://doi.org/10.1186/s12870-020-2326-5>
- Burns, R. M. B. H. H. (Ed.). (1990). *Silvics of North America*. U.S. Department of Agriculture, Forest Service.
- Cao, Y., He, L., Song, F., Li, C., Ji, Q., Liu, J., Peng, G., Li, B., Zeng, F., & Zhan, Y. (2023). Physiological and gene expression response of interspecific hybrids of *Fraxinus mandshurica* × *Fraxinus americana* to MJ or SNP under drought. *Forests*, 14, 1277. <https://doi.org/10.3390/f14061277>
- Chen, J., Zhao, J., Erickson, D. L., Xia, N., & Kress, W. J. (2015). Testing DNA barcodes in closely related species of *curcuma* (Zingiberaceae) from Myanmar and China. *Mol Ecol Resour.*, 15(2), 337–348. <https://doi.org/10.1111/1755-0998.12319>
- Coker, T. L. R., Rozsypálek, J., Edwards, A., Harwood, T. P., Butfoy, L., & Buggs, R. J. A. (2019). Estimating mortality rates of European ash (*Fraxinus excelsior*) under the ash dieback (*Hymenoscyphus fraxineus*) epidemic. *Plants, People, Planet*, 1, 48–58. <https://doi.org/10.1002/ppp3.11>
- Davidson, C. G. (1999). ‘Northern treasure’ and ‘Northern gem’ hybrid ash. *HortScience*, 34, 151–152. <https://doi.org/10.21273/HORTSCI.34.1.151>
- Davidson CG Ronald WG 2001 Hybrid ash tree named ‘Northern Treasure’. *US Patent App.* 09/233,440
- Dobrowolska, D., Hein, S., Oosterbaan, A., Wagner, S., Clark, J., & Skovsgaard, J. P. (2011). A review of European ash (*Fraxinus excelsior* L.): Implications for silviculture. *Forestry*, 84, 133–148. <https://doi.org/10.1093/forestry/cpr001>
- Dolezel, J., Greilhuber, J., & Suda, J. (2007). Estimation of nuclear DNA content in plants using flow cytometry. *Nature Protocols*, 2, 2233–2244. <https://doi.org/10.1038/nprot.2007.310>
- Drenkhan, R., Agan, A., Palm, K., Rosenvald, R., Jürisoo, L., Maaten, T., Padari, A., & Drenkhan, T. (2017). Overview of ash and ash dieback in Estonia. In R. Vasaitis & R. Enderle (Eds.), *Dieback of European ash (Fraxinus spp.): Consequences and guidelines for sustainable management* (pp. 115–124). Swedish University of Agricultural Sciences.
- Edgar, R. C. (2004). MUSCLE: Multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research*, 32, 1792–1797. <https://doi.org/10.1093/nar/gkh340>
- Ensslin, A., & Godefroid, S. (2019). How the cultivation of wild plants in botanic gardens can change their genetic and phenotypic status and what this means for their conservation value. *Sibbaldia*, 17, 51–70.
- Fernández-Manjarrés, J. F., Gerard, P. R., Dufour, J., Raquin, C., & Frascaria-Lacoste, N. (2006). Differential patterns of morphological and molecular hybridization between *Fraxinus excelsior* L. and *Fraxinus angustifolia* Vahl (Oleaceae) in eastern and western France. *Molecular Ecology*, 15, 3245–3257. <https://doi.org/10.1111/j.1365-294X.2006.02975.x>
- Gamborg, O. L., Miller, R., & Ojima, K. (1968). Nutrient requirements of suspension cultures of soybean root cells. *Experimental Cell Research*, 50(1), 151–158.
- Gao, T., Yao, H., Song, J., Zhu, Y., Liu, C., & Chen, S. (2010). Evaluating the feasibility of using candidate DNA barcodes in discriminating species of the large Asteraceae family. *BMC Evolutionary Biology*, 10, 324. <https://doi.org/10.1186/1471-2148-10-324>
- Gross, A., Holdenrieder, O., Pautasso, M., Quelo, V., & Sieber, T. N. (2014). *Hymenoscyphus pseudoalbidus*, the causal agent of European ash dieback. *Molecular Plant Pathology*, 15, 5–21. <https://doi.org/10.1111/mpp.12073>
- He, L., Xu, Y., Zeng, F., Tian, H., Xiao, Y., Liu, H., Yu, L., & Zhan, Y. (2021). Establishment of a micropropagation supporting technology for the *Fraxinus mandshurica* × *Fraxinus sogdiana*. *In Vitro Cellular & Developmental Biology - Plant*, 57, 307–318. <https://doi.org/10.1007/s11627-021-10157-5>
- He, L., Yan, J., Lu, H., Su, W., Gao, S., Wang, Y., Zhan, Y., & Zeng, F. (2023). Phenotypic differences of leaves and transcriptome analysis of *Fraxinus mandshurica* × *Fraxinus sogdiana* F1 variety. *Forests, Trees and Livelihoods*, 14, 1554. <https://doi.org/10.3390/f14081554>
- He, L., Zeng, F., He, Z., Li, Y., Li, S., & Zhan, Y. (2019). Interspecific hybridizations of *Fraxinus* L. (*F. mandshurica* × *F. americana* and *F. mandshurica* × *F. velutina*) and heterosis analysis and selection of F1 progenies. *Canadian Journal of Forest Research*, 49, 1265–1276. <https://doi.org/10.1139/cjfr-2018-0460>
- Henry, A. (1914). The artificial production of vigorous trees. *Department of Agriculture and Technical Instruction for Ireland Journal*, XV, 34–52.
- Herms, D. A., & McCullough, D. G. (2014). Emerald ash borer invasion of North America: History, biology, ecology, impacts, and management. *Annual Review of Entomology*, 59, 13–30. <https://doi.org/10.1146/annurev-ento-011613-162051>
- Heuert, M., Carnevale, S., Fineschi, S., Sebastiani, F., Hausman, J. F., Paule, L., & Vendramin, G. G. (2006). Chloroplast DNA phylogeography of European ashes, *Fraxinus* sp. (Oleaceae): Roles of hybridization and life history traits. *Molecular Ecology*, 15, 2131–2140. <https://doi.org/10.1111/j.1365-294X.2006.02897.x>

- Hinsinger, D. D., Basak, J., Gaudeul, M., Cruaud, C., Bertolino, P., Frascaria-Lacoste, N., & Bousquet, J. (2013). The phylogeny and biogeographic history of ashes (*Fraxinus*, Oleaceae) highlight the roles of migration and vicariance in the diversification of temperate trees. *PLoS One*, 8, e80431. <https://doi.org/10.1371/journal.pone.0080431>
- Hinsinger, D. D., Gaudeul, M., Couloux, A., Bousquet, J., & Frascaria-Lacoste, N. (2014). The phylogeography of Eurasian *Fraxinus* species reveals ancient transcontinental reticulation. *Molecular Phylogenetics and Evolution*, 77, 223–237. <https://doi.org/10.1016/j.ympev.2014.04.021>
- Jeandroz, S., Roy, A., & Bousquet, J. (1997). Phylogeny and phylogeography of the circumpolar genus *Fraxinus* (Oleaceae) based on internal transcribed spacer sequences of nuclear ribosomal DNA. *Molecular Phylogenetics and Evolution*, 7, 241–251. <https://doi.org/10.1006/mpev.1996.0393>
- Jepson, P. R., & Arakelyan, I. (2017). Developing publicly acceptable tree health policy: Public perceptions of tree-breeding solutions to ash dieback among interested publics in the UK. *Forest Policy and Economics*, 80, 167–177. <https://doi.org/10.1016/j.forpol.2017.03.002>
- Johnson, L. P. V., & Heimbürger, C. (1946). Preliminary report on interspecific hybridization in forest trees. *Canadian Journal of Research*, 24c, 308–312.
- Kelly, L. J., Coates, S. E. R., Lee, S. J., Rossiter, S. J., & Buggs, R. J. A. (2025). Molecular convergence analyses identify candidate genes for low susceptibility to the ash dieback pathogen. *bioRxiv*. <https://doi.org/10.1101/2025.02.08.637240>
- Kelly, L. J., Plumb, W. J., Carey, D. W., Mason, M. E., Cooper, E. D., Crowther, W., Whittemore, A. T., Rossiter, S. J., Koch, J. L., & Buggs, R. J. A. (2020). Convergent molecular evolution among ash species resistant to the emerald ash borer. *Nature Ecology & Evolution*, 4, 1116–1128. <https://doi.org/10.1038/s41559-020-1209-3>
- Kim, C., Kim, D.-K., Sun, H., & Kim, J.-H. (2022). Phylogenetic relationship, biogeography, and conservation genetics of endangered *Fraxinus chii-sanensis* (Oleaceae), endemic to South Korea. *Plant Diversity*, 44, 170–180. <https://doi.org/10.1016/j.pld.2021.06.004>
- Knox, R. B., Gaget, M., & Dumas, C. (1987). Mentor pollen techniques. In G. H. Bourne, K. W. Jeon, & M. Friedlander (Eds.), *International review of cytology* (pp. 315–332). Academic Press.
- Koch, J. L. (2025). Proof of a genetic basis for EAB-resistance: An update on breeding research. In K. A. McManus (Ed.), *Proceedings, 29th USDA interagency research forum on invasive species*. U.S. Department of Agriculture, Forest Service, Forest Health Technology Enterprise Team.
- Koch, J. L., Carey, D. W., Knight, K. S., Poland, T., Herms, D. A., & Mason, M. E. (2012). Breeding strategies for the development of emerald ash borer-resistant North American ash. In *Proceedings of the 4th international workshop on the genetics of host-parasite interactions in forestry: Disease and insect resistance in forest trees* (pp. 235–239). USDA Forest Service, Pacific Southwest Research Station Albany.
- Koch, J. L., Carey, D. W., & Mason, M. E. (2007). Development of novel ash hybrids to introgress resistance to emerald ash borer into north American ash species. In V. Mastro, D. Lance, R. Reardon, & G. Parra (Eds.), *Emerald ash borer research and development meeting* (pp. 36–37). U.S. Department of Agriculture, Forest Service, Forest Health Technology Enterprise Team.
- Koch, J. L., Carey, D. W., Mason, M. E., & Nurul Islam-Faridi, M. (2010). Overcoming obstacles to interspecies hybridization of ash. In C. H. Michler & M. D. Ginzler (Eds.), *General technical report - proceedings. Proceedings of symposium on ash in North America; 2010 March* (pp. 41–44). U.S. Department of Agriculture, Forest Service.
- Koch, J. L., Carey, D. W., Mason, M. E., Poland, T. M., & Knight, K. S. (2015). Intraspecific variation in *Fraxinus pennsylvanica* responses to emerald ash borer (*Agrilus planipennis*). *New Forests*, 46, 995–1011. <https://doi.org/10.1007/s11056-015-9494-4>
- Kowalski, T., Bilański, P., & Holdenrieder, O. (2015). Virulence of *Hymenoscyphus albidus* and *H. Fraxineus* on *Fraxinus excelsior* and *F. Pennsylvanica*. *PLoS One*, 10, e0141592. <https://doi.org/10.1371/journal.pone.0141592>
- Kozlov, A. M., Darriba, D., Flouri, T., Morel, B., & Stamatakis, A. (2019). RAxML-NG: A fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics*, 35, 4453–4455. <https://doi.org/10.1093/bioinformatics/btz305>
- Lévesque, M., Bustamante Eduardo, J. I., & Queloz, V. (2023). Potential alternative tree species to *Fraxinus excelsior* in European forests. *Frontiers in Forests and Global Change*, 6, 1048971. <https://doi.org/10.3389/ffgc.2023.1048971>
- Lingelsheim, A. (1907). Vorarbeiten zu einer Monographie der Gattung *Fraxinus*. *Botanische Jahrbücher Für Systematik, Pflanzengeschichte Und Pflanzengeographie*, 40, 185–223.
- Liu, J. N., Yan, L., Chai, Z., Liang, Q., Dong, Y., Wang, C., Li, X., Li, C., Mu, Y., Gong, A., et al. (2024). Pan-genome analyses of 11 *Fraxinus* species provide insights into salt adaptation in ash trees. *Plant Communications*, 0, 101137.
- Liu, X., Xing, H., Kong, F., Zhang, K., Cao, Y., Guo, X., Li, Q., Wang, J., Jing, T., Zhan, Y., & Qi, F. (2025). Molecular identification of F₁ hybrids of *Fraxinus mandshurica* × *Fraxinus chinensis* using SSR markers. *Gene*, 959, 149507. <https://doi.org/10.1016/j.gene.2025.149507>
- Lloyd, G., & McCown, B. (1980). Commercially-feasible micropropagation of mountain laurel, *Kalmia latifolia*, by use of shoot-tip culture. *Combined Proceedings, International Plant Propagators' Society*, 30, 421–427.
- Loureiro, J., Rodriguez, E., Dolezel, J., & Santos, C. (2007). Two new nuclear isolation buffers for plant DNA flow cytometry: A test with 37 species. *Annals of Botany*, 100, 875–888. <https://doi.org/10.1093/aob/mcm152>
- Marzano, M., Woodcock, P., & Quine, C. P. (2019). Dealing with dieback: Forest manager attitudes towards developing resistant ash trees in the United Kingdom. *Forestry: An International Journal of Forest Research*, 92, 554–567. <https://doi.org/10.1093/forestry/cpz022>
- Maunder, M., Hughes, C., Hawkins, J. A., & Culham, A. (2004). Hybridization in ex situ plant collections: conservation concerns, liabilities, and opportunities. In M. Maunder & EOG. (Eds.), *Ex situ plant conservation* (pp. 325–363). Island Press.
- Miller, G. N. (1955). *The genus Fraxinus, the ashes, in North America, north of Mexico* (Vol. 355) (pp. 1–64). Cornell University Agricultural Experimental Station Memoir.
- Mitchell, R. J., Beaton, J. K., Bellamy, P. E., Broome, A., Chetcuti, J., Eaton, S., Ellis, C. J., Gimona, A., Harmer, R., Hester, A. J., Beaton, J. K., Bellamy, P. E., Broome, A., Chetcuti, J., Eaton, S., Ellis, C. J., Gimona, A., Harmer, R., Hester, A. J., ... Woodward, S. (2014). Ash dieback in the UK: A review of the ecological and conservation implications and potential management options. *Biological Conservation*, 175, 95–109. <https://doi.org/10.1016/j.biocon.2014.04.019>
- Nielsen, L. R., McKinney, L. V., Hietala, A. M., & Kjær, E. D. (2016). The susceptibility of Asian, European and north American *Fraxinus* species to the ash dieback pathogen *Hymenoscyphus fraxineus* reflects their phylogenetic history. *European Journal of Forest Research*, 136(1), 1–73. <https://doi.org/10.1007/s10342-016-1009-0>
- Nikolaev, E. V. (1981). The genus *Fraxinus* (Oleaceae) in the flora of the USSR. *Botanicheskii Zhurnal*, 66, 1419–1432.
- Obermayer, R., Leitch, I. J., Hanson, L., & Bennett, M. D. (2002). Nuclear DNA c-values in 30 species double the familial representation in pteridophytes. *Annals of Botany*, 90, 209–217. <https://doi.org/10.1093/aob/mcf167>
- Orlova-Bienkowskaja, M. J., & Bieńkowski, A. O. (2016). The life cycle of the emerald ash borer *Agrilus planipennis* in European Russia and comparisons with its life cycles in Asia and North America. *Agricultural and Forest Entomology*, 18(2), 182–188. <https://doi.org/10.1111/afe.12140>

- Orlova-Bienkowskaja, M. J., & Bieńkowski, A. O. (2018). Modeling long-distance dispersal of emerald ash borer in European Russia and prognosis of spread of this pest to neighboring countries within next 5 years. *Ecology and Evolution*, 8, 9295–9304. <https://doi.org/10.1002/ece3.4437>
- Pautasso, M., Aas, G., Queloz, V., & Holdenrieder, O. (2013). European ash (*Fraxinus excelsior*) dieback—A conservation biology challenge. *Biological Conservation*, 158, 37–49. <https://doi.org/10.1016/j.biocon.2012.08.026>
- Petit, R. J., Bodénès, C., Ducouso, A., Roussel, G., & Kremer, A. (2004). Hybridization as a mechanism of invasion in oaks. *New Phytologist*, 161, 151–164. <https://doi.org/10.1046/j.1469-8137.2003.00944.x>
- Plumb, W. J., Coker, T. L. R., Stocks, J. J., Woodcock, P., Quine, C. P., Nemesio-Gorriz, M., Douglas, G. C., Kelly, L. J., & Buggs, R. J. A. (2019). The viability of a breeding programme for ash in the British Isles in the face of ash dieback. *Plants People Planet*, 58, 227.
- Procter, J. B., Carstairs, G. M., Soares, B., Mourão, K., Ofoegbu, T. C., Barton, D., Lui, L., Menard, A., Sherstnev, N., Roldan-Martinez, D., Carstairs, G. M., Soares, B., Mourão, K., Ofoegbu, T. C., Barton, D., Lui, L., Menard, A., Sherstnev, N., Roldan-Martinez, D., ... Barton, G. J. (2021). Alignment of biological sequences with Jalview. *Methods in Molecular Biology*, 2231, 203–224. https://doi.org/10.1007/978-1-0716-1036-7_13
- Raquin, C., Jung-Muller, B., Dufour, J., & Frascaria-Lacoste, N. (2002). Rapid seedling obtaining from European ash species *Fraxinus excelsior* (L.) and *Fraxinus angustifolia* (Vahl.). *Annals of Forest Science*, 59, 219–224.
- Ren, F. M., Wang, Y. W., Xu, Z. C., Li, Y., Xin, T. Y., Zhou, J. G., Qi, Y. D., Wei, X. P., Yao, H., & Song, J. Y. (2019). DNA barcoding of *corydalis*, the most taxonomically complicated genus of Papaveraceae. *Ecology and Evolution*, 9(4), 1934–1945. <https://doi.org/10.1002/ece3.4886>
- Rigsby, C. M., Herms, D. A., Bonello, P., & Cipollini, D. (2016). Higher activities of defense-associated enzymes may contribute to greater resistance of Manchurian ash to Emerald ash borer than a closely related and susceptible congener. *Journal of Chemical Ecology*, 42, 782–792. <https://doi.org/10.1007/s10886-016-0736-5>
- Santini, A., Fagnani, A., Ferrini, F., Ghelardini, L., & Mittempergher, L. (2007). ‘Fiorente’ and ‘Arno’ elm trees. *HortScience*, 42, 712–714. <https://doi.org/10.21273/HORTSCI.42.3.712>
- Sela, I., Ashkenazy, H., Katoh, K., & Pupko, T. (2015). Guidance2: Accurate detection of unreliable alignment regions accounting for the uncertainty of multiple parameters. *Nucleic Acids Research*, 43, W7–W14. <https://doi.org/10.1093/nar/gkv318>
- Sharma, D. R., Kaur, R., & Kumar, K. (1996). Embryo rescue in plants—A review. *Euphytica*, 89, 325–337. <https://doi.org/10.1007/BF00022289>
- Siljak-Yakovlev, S., Temunović, M., Robin, O., Raquin, C., & Frascaria-Lacoste, N. (2014). Molecular-cytogenetic studies of ribosomal RNA genes and heterochromatin in three European *Fraxinus* species. *Tree Genetics and Genomes*, 10, 231–239. <https://doi.org/10.1007/s11295-013-0654-4>
- Smalley, E. B., & Guries, R. P. (1993). Breeding elms for resistance to Dutch elm disease. *Annual Review of Phytopathology*, 31, 325–354. <https://doi.org/10.1146/annurev.py.31.090193.001545>
- Sollars ESA, Harper AL, Kelly LJ, Sambles CM, Ramirez-Gonzalez RH, Swarbreck D, Kaithakottil G, Cooper ED, Uauy C, Havlickova L, Harper AL, Kelly LJ, Sambles CM, Ramirez-Gonzalez RH, Swarbreck D, Kaithakottil G, Cooper ED, Uauy C, Havlickova L, Kaithakottil G, Cooper ED, Uauy C, Havlickova L, Worswick G, Studholme DJ, Zohren J, Salmon DL, Clavijo BJ, Li Y, He Z, Fellgett A, McKinney LV, Nielsen LR, Douglas GC, Kjaer ED, Downie JA, Boshier D, Lee S, Clark J, Grant M, Bancroft I, Caccamo M, Buggs RJA 2017. Genome sequence and genetic diversity of European ash trees. *Nature* 541: 212–216, <https://doi.org/10.1038/nature20786>
- Thomasset, M., Fernandez-Manjarrés, J. F., Douglas, G. C., Frascaria-Lacoste, N., Raquin, C., & Hodkinson, T. R. (2011). Molecular and morphological characterization of reciprocal F1 hybrid ash (*Fraxinus excelsior* × *Fraxinus angustifolia*, Oleaceae) and parental species reveals asymmetric character inheritance. *International Journal of Plant Sciences*, 172, 423–433.
- Wallander, E. (2008). Systematics of *Fraxinus* (Oleaceae) and evolution of dioecy. *Plant Systematics and Evolution*, 273, 25–49. <https://doi.org/10.1007/s00606-008-0005-3>
- Wallander, E. (2012). Systematics and floral evolution in *Fraxinus* (Oleaceae). *Belgische Dendrologie Belge*, 2013, 39–58.
- Wenzig, T. (1883). Die Gattung *Fraxinus* Tourn. neu bearbeitet. *Botanische Jahrbücher für Systematik, Pflanzengeschichte Und Pflanzengeographie*, 4, 165–188.
- Wesmael, A. (1892). Monographie des espèces du genre *Fraxinus*. *Bulletin de la Société Royale de Botanique de Belgique = Bulletin Van de Koninklijke Belgische Botanische Vereniging*, 31, 69–117.
- Whittemore, A. T., Campbell, J. J. N., Xia, Z.-L., Carlson, C. H., Atha, D., & Olsen, R. T. (2018). Ploidy variation in *Fraxinus* L. (Oleaceae) of eastern North America: Genome size diversity and taxonomy in a suddenly endangered genus. *International Journal of Plant Sciences*, 179, 377–389. <https://doi.org/10.1086/696688>
- Wright, J. W. (1953). *Summary of tree-breeding experiments by the north-eastern Forest Experiment Station 1947-1950*. Northeastern Forest Experiment Station.
- Zeng, F.-S., Li, L.-L., Liang, N.-S., Wang, X., Li, X., & Zhan, Y.-G. (2015). Salt tolerance and alterations in cytosine methylation in the interspecific hybrids of *Fraxinus velutina* and *Fraxinus mandshurica*. *Euphytica*, 205, 721–737. <https://doi.org/10.1007/s10681-015-1432-1>

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Plumb, W. J., Kelly, L. J., Mullender, J., Powell, R. F., Csiba, L., Nemesio-Gorriz, M., Carey, D., Mason, M. E., Crowther, W., Koch, J., Douglas, G. C., & Buggs, R. J. A. (2025). Preliminary genetic barcodes for ash (*Fraxinus*) species and generation of new wide hybrids. *Plants, People, Planet*, 1–14. <https://doi.org/10.1002/ppp3.70069>